How to use the
BTNX Rapid Tests

BEFORE

Blow your nose and wash your hands with soap

1 SET UP



Place the tube in the holder

Fill the tube with the test liquid

2 COLLECTION

A



Peel down and remove nasal swab

B



Insert swab to reach back of throat

Rub nasal swab against upper part of back of throat. **Avoid tongue, teeth and gums**



C

Insert nasal swab **1/2 inch** from edge of nostril. Roll **5 times in each nostril** and keep the nasal swab in place for **10 seconds** (left & right).

3 PREPARATION

Insert the swab in the tube liquid and rotate it **10 times**



Place the tube and swab in the holder and leave it there for **2 minutes**

Remove the swab by pinching the tube



Close the tube with the stopper cap

4 TESTING

Place **3 drops** in the rapid response cartridge and wait **15 minutes**

(Do not touch it during the waiting time)

Here is an example of a positive test, a negative test and an invalid test



NEGATIVE POSITIVE INVALID

Redo the nasal swab with a new kit.

* If your rapid test is **NEGATIVE**, you must **do a second test 48 hours later**

If you are POSITIVE, refer to the MSSS guidelines. For any questions, contact line 811.

ᓄᓇᕕᒻᒥ ᐃᓗᓯᓕᕆᓂᕐᒧᑦ ᑲᑎᒪᔨᑦ
RÉGIE RÉGIONALE DE LA NUNAVIK REGIONAL SANTÉ ET DES SERVICES BOARD OF HEALTH SOCIAUX DU NUNAVIK AND SOCIAL SERVICES